

Package ‘swfdr’

October 16, 2019

Title Science-wise false discovery rate and proportion of true null hypotheses estimation

Version 1.10.0

Author Jeffrey T. Leek, Leah Jager, Simina M. Boca

Maintainer Simina M. Boca <smb310@georgetown.edu>, Jeffrey T. Leek
<jtleek@gmail.com>

Description This package allows users to estimate the science-wise false discovery rate from Jager and Leek, ``Empirical estimates suggest most published medical research is true," 2013, Biostatistics, using an EM approach due to the presence of rounding and censoring. It also allows users to estimate the proportion of true null hypotheses in the presence of covariates, using a regression framework, as per Boca and Leek, ``A regression framework for the proportion of true null hypotheses," 2015, bioRxiv preprint.

Depends R (>= 3.4)

Imports stats4, ggplot2, reshape2, stats, dplyr

License GPL (>= 3)

Encoding UTF-8

LazyData true

RoxygenNote 6.0.1

Suggests BiocStyle, knitr, rmarkdown

VignetteBuilder knitr

biocViews MultipleComparison, StatisticalMethod, Software

git_url <https://git.bioconductor.org/packages/swfdr>

git_branch RELEASE_3_9

git_last_commit 35610f9

git_last_commit_date 2019-05-02

Date/Publication 2019-10-15

R topics documented:

BMI_GIANT_GWAS_sample	2
calculateSwfdr	3
journals_pVals	4
lm_pi0	4

Index

6

BMI_GIANT_GWAS_sample *Subset of SNPs from meta-analysis of BMI GWAS study.*

Description

A dataset containing 50,000 SNPs and results for their associations with BMI.

Usage

```
data(BMI_GIANT_GWAS_sample)
```

Format

A data frame with 50,000 rows and 9 variables:

SNP ID for SNP (single nucleotide polymorphism)

A1 Allele 1 for SNP

A2 Allele 2 for SNP

Freq_MAF_Hapmap Frequency of minor allele (MAF) in Hapmap project

b Estimated beta for association between SNP and BMI

se Estimated standard error (se) for association between SNP and BMI

p P-value for association between SNP and BMI

N Total sample size considered for association of SNP and BMI

Freq_MAF_Int_Hapmap Three approximately equal intervals for the Hapmap MAFs

Value

Object of class `tbl_df`, `tbl`, `data.frame`.

Source

https://www.broadinstitute.org/collaboration/giant/index.php/GIANT_consortium_data_files#GWAS_Anthropometric_2015_BMI

calculateSwfdr	<i>Calculate the science-wise FDR (swfdr)</i>
----------------	---

Description

Calculate the science-wise FDR (swfdr)

Usage

```
calculateSwfdr(pValues, truncated, rounded, pi0 = 0.5, alpha = 1,
               beta = 50, numEmIterations = 100)
```

Arguments

pValues	Numerical vector of p-values
truncated	Vector of 0s and 1s with indices corresponding to those in pValues; 1 indicates that the p-values is truncated, 0 that it is not truncated
rounded	Vector of 0s and 1s with indices corresponding to those in pValues; 1 indicates that the p-values is rounded, 0 that it is not rounded
pi0	Initial prior probability that a hypothesis is null (default is 0.5)
alpha	Initial value of parameter alpha from Beta(alpha, beta) true positive distribution (default is 1)
beta	Initial value of parameter beta from Beta(alpha, beta) true positive distribution (default is 50)
numEmIterations	The number of EM iterations (default is 100)

Value

pi0	Final value of prior probability - estimated from EM - that a hypothesis is null, i.e. estimated swfdr
alpha	Final value of parameter alpha - estimated from EM - from Beta(alpha, beta) true positive distribution
beta	Final value of parameter beta - estimated from EM - from Beta(alpha, beta) true positive distribution
z	Vector of expected values of the indicator of whether the p-value is null or not - estimated from EM - for the non-rounded p-values (values of NA represent the rounded p-values)
n0	Expected number of rounded null p-values - estimated from EM - between certain cutpoints (0.005, 0.015, 0.025, 0.035, 0.045, 0.05)
n	Number of rounded p-values between certain cutpoints (0.005, 0.015, 0.025, 0.035, 0.045, 0.05)

Examples

```
pVals <- runif(100)
tt <- rr <- rep(0, 100)
resSwfdr <- calculateSwfdr(pValues = pVals, truncated = tt, rounded = rr, numEmIterations=100)
```

journals_pVals	<i>P-values from abstracts from articles in 5 biomedical journals (American Journal of Epidemiology, BMJ, JAMA, Lancet, New England Journal of Medicine), over 11 years (2000-2010).</i>
----------------	--

Description

A dataset containing 15,653 p-values.

Usage

```
journals_pVals
```

Format

A `tbl` data frame with 15,653 rows and 5 variables:

pvalue P-value

pvalueTruncated Equals to 1 if the p-value is truncated, 0 otherwise

pubmedID Pubmed ID of the article

year Year of publication

journal Journal

Value

Object of class `tbl_df`, `tbl`, `data.frame`.

Source

Code for extracting p-values at: [inst/script/getPvalues.R](#)

lm_pi0

Estimate pi0(x)

Description

Estimate $\pi_0(x)$

Usage

```
lm_pi0(pValues, lambda = seq(0.05, 0.95, 0.05), X, type = "logistic",
smooth.df = 3, threshold = TRUE)
```

Arguments

pValues	Numerical vector of p-values
lambda	Numerical vector of thresholds. Must be in [0,1).
X	Design matrix (one test per row, one variable per column). Do not include the intercept.
type	Type of regression, "logistic" or "linear." Default is logistic.
smooth.df	Number of degrees of freedom when estimating pi0(x) with a smoother.
threshold	If TRUE (default), all estimates are thresholded at 0 and 1, if FALSE, none of them are.

Value

- pi0 Numerical vector of smoothed estimate of $\pi_0(x)$. The length is the number of rows in X.
 pi0.lambda Numerical matrix of estimated $\pi_0(x)$ for each value of lambda. The number of columns is the number of tests, the number of rows is the length of lambda.
 lambda Vector of the values of lambda used in calculating pi0.lambda
 pi0.smooth Matrix of fitted values from the smoother fit to the $\pi_0(x)$ estimates at each value of lambda (same number of rows and columns as pi0.lambda)

Examples

```
X <- seq(-1,2,length=1000) ##covariate
pi0 <- 1/4*X + 1/2 ##probability of being null
nullI <- rbinom(1000,prob=pi0,size=1)> 0 ##generate null/alternative p-values
pValues <- rep(NA,1000) ##vector of p-values
pValues[nullI] <- runif(sum(nullI)) ##null from U(0,1)
pValues[!nullI] <- rbeta(sum(!nullI),1,2) ##alternative from Beta
pi0x <- lm_pi0(pValues=pValues, X=X, smooth=df=3)
```

Index

*Topic **datasets**

BMI_GIANT_GWAS_sample, [2](#)

journals_pVals, [4](#)

BMI_GIANT_GWAS_sample, [2](#)

calculateSwfdr, [3](#)

journals_pVals, [4](#)

lm_pi0, [4](#)